

# Math 126 Midterm 1 Problems

## P.1) True/False

A.) The method of characteristics guarantees a solution to

$$\left\{ \begin{array}{l} \partial_t u + u \partial_x u = 0 \end{array} \right.$$

$$\left\{ \begin{array}{ll} u(0, x) = \begin{cases} 1 & x \leq 0 \\ 0 & x \geq 1 \end{cases} & x \in [0, 1] \end{array} \right.$$

for all time.

False - HW 1 Problem 4

B.) If you lived in 2-Dimensions and someone clapped, you would hear a sharp noise followed by silence.

False - Interpreting Huygen's Principle - See lecture 8 or figure 4.14 (page 66) in the text

C.) Let  $U \subseteq \mathbb{R}^n$  be a bounded domain with piecewise- $C^1$  boundary.

A solution  $u \in C^2([0, \infty) \times \bar{U})$  to

$$\left\{ \begin{array}{l} \frac{\partial^2 u}{\partial t^2} - c^2 \Delta u = f \\ u(0, x) = g \\ \frac{\partial u}{\partial t}(0, x) = h \\ u|_{\partial U} = 0 \end{array} \right.$$

is unique for  $f \in C^0$ ,  $g \in C^2$ ,  $h \in C^1$ .

True - Corollary 4.13

D.) Eigenvalues of the Laplace operator ( $\lambda$  solving  $-\Delta \phi = \lambda \phi$ )

correspond to frequencies we observe when a string

vibrates as governed by the wave equation with fixed ends.

True - Section 5.2 of the text or  
lecture 9

E.) If  $U \subseteq \mathbb{R}^n$  is a bold-domain with piecewise- $C^1$  bdry,

$$\int_U u \Delta u dx = \int_{\partial U} \frac{\partial u}{\partial \eta} dS + \int_U |\nabla u|^2 dx$$

False -  $\int_U u \Delta u dx = \int_{\partial U} u \frac{\partial u}{\partial \eta} dS - \int_U |\nabla u|^2 dx$ .

P2) Let  $u(t, x)$  solve  $\frac{\partial u}{\partial t} + \frac{\partial u}{\partial x} = 1$  in the first quadrant  
 $Q_1 = \{(t, x) \mid t \geq 0, x \geq 0\}$ . Suppose  
 $\begin{cases} u(0, x) = x & x \geq 0 \\ u(t, 0) = t & t \geq 0 \end{cases}$

Find a formula for  $u \in C^1(Q_1)$  using the method of characteristics.

Solution: HW 1 P283

The characteristic ODE is  $\dot{x}(t) = 1$  with  $x(0) = x_0$   
or  $x(t_0) = 0$  depending on boundary data.

Then,  $x(t) = x_0 + t$  for  $x_0 \geq 0$   
or  $x(t) = t - t_0$  for  $t_0 \geq 0$

Next,  $\frac{D u}{D t} = 1$  gives  $u(t, x(t)) = \begin{cases} t + u(0, x_0) & x \geq t \\ t - t_0 + u(t_0, 0) & x \leq t \end{cases}$

$$\text{So } u(t, x) = \begin{cases} t + x - t & x \geq t \\ t - t_0 + t_0 & x \leq t \end{cases} = \begin{cases} x & x \geq t \\ t & x \leq t \end{cases}$$


P.3) Assume that  $u(t, x) \in C^2([0, \infty) \times \mathbb{R}_x)$  solves

$$\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = 0.$$

A.) Let  $u(0, 1) = 2$ ,  $u(1, 2) = 4$ , and  $u(1, 0) = 3$ .

What is  $u(2, 1)$ .

Solution: As in HW 2 P1, these points form a parallelogram whose sides are characteristics



By the formula from that problem,  $u(2, 1) = u(1, 2) + u(1, 0) - u(0, 1)$   
 $= 7 - 2 = 5$ .

B.) Let  $v$  solve  $\frac{\partial^2 v}{\partial t^2} - \frac{\partial^2 v}{\partial x^2} = 1$  and  $v(0, x) = u(0, x)$ ,

$\partial_t v(0, x) = \partial_t u(0, x)$ . What is  $v(2, 1)$ ?

Solution:


Since  $v-u$  solves

$$\left\{ \begin{array}{l} \frac{\partial^2 w}{\partial t^2} - \frac{\partial^2 w}{\partial x^2} = 1 \\ w(0, x) = 0 \\ \partial_t w(0, x) = 0 \end{array} \right.$$

$$v(t, x) = u(t, x) + w(t, x) = u(t, x) + \frac{1}{2} \int_{D_{t,x}} 1 \, ds dt.$$

$$\begin{aligned} \text{Then, } v(2, 1) &= 5 + \frac{1}{2} \cdot \text{Area}(\{(t, x) : 1 - (2-t) \leq x \leq 1 + (2-t)\}) \\ &= 5 + \frac{1}{2}(4) = 7. \end{aligned}$$

C.) What is the range of influence of  $(z, 1)$ ?



$$T^+_{(z,1)} = \{(t, x) \mid z-t \leq x \leq t-1\}$$